

TRANSITIVE GROUP ACTIONS: (IM)PRIMITIVITY AND SEMIREGULAR SUBGROUPS

István Kovács^{a,1, 2}, Aleksander Malnič^{b,1}, Dragan Marušič^{a, b, 1, *}, Štefko Miklavic^{a, b, 1}

^a*University of Primorska, Cankarjeva 5, 6000 Koper, Slovenia and*

^b*IMFM, University of Ljubljana, Jadranska 19, 1111 Ljubljana, Slovenia*

Abstract

The following problem is considered: if H is a semiregular abelian subgroup of a transitive permutation group G acting on a finite set X , find conditions for (non)existence of G -invariant partitions of X and give their description in terms of orbits of H . Conditions and descriptions presented in this paper are derived by studying spectral properties of the associated G -invariant digraphs arising as regular abelian covers. As an essential tool, irreducible complex characters defined on H , are used.

Questions of this kind arise naturally when classifying combinatorial objects which enjoy a certain degree of symmetry. The techniques of this paper allow one to consider such problems without referring to, as it often is the case, the classification of finite simple groups. As an illustration, a new and short proof of an old result of Frucht, Graver and Watkins (*Proc. Camb. Phil. Soc.*, **70** (1971), 211-218) classifying edge-transitive generalized Petersen graphs, is given.

1 Introductory and historic remarks

When investigating the structure of graphs admitting transitive actions (or possibly intransitive actions with a small number of orbits) of subgroups of automorphisms one often relies on results which are purely group-theoretic in nature. On the other hand, graph-theoretic language can often provide useful insights in the study of various problems in permutation groups.

A particular case of such an interplay of group-theoretic and graph-theoretic concepts is dealt with in this paper, and is motivated by the following situation.

¹Supported in part by “Ministrstvo za visoko šolstvo, znanost in tehnologijo Slovenije”, program no. P1-0285, and the Slovenian-Hungarian Intergovernmental and Technological Cooperation SLO-9/2005.

²Supported in part by the grant OTKA T-043758.

* Corresponding author e-mail: dragan.marusic@guest.arnes.si

Given a *bicirculant*, that is, a graph admitting a cyclic subgroup with two orbits of equal size, there are two essentially different possibilities forcing such a graph to be vertex-transitive. Either there exists an automorphism swapping setwise the two orbits, called hereafter a *swap*, or there exists an automorphism mixing the vertices of the two orbits, called hereafter a *mixer*. For example, the class of generalized Petersen graphs $GP(n, s)$ contains examples of each of the four possibilities that may occur with this respect: first, $GP(7, 2)$ is not vertex-transitive and hence without swaps and mixers, second, the prism $GP(3, 1)$ has swaps, but no mixers, third, the dodecahedron $GP(10, 2)$ has mixers, but no swaps, and finally, the cube $GP(4, 1)$ has both swaps and mixers. It seems therefore natural to seek for (non)existence conditions for these two kinds of automorphisms in bicirculants.

When attempting to answer questions of this nature, it is often useful to choose a more general framework. We do this here by considering arbitrary transitive permutation groups (not necessarily automorphism groups of graphs) containing semiregular abelian subgroups (not necessarily cyclic subgroups with two orbits of equal size). Recall that a nontrivial permutation, and more generally a nontrivial permutation group, is *semiregular* if all of its orbits have equal size. We note that asking of a transitive permutation group to contain a semiregular abelian subgroup is not that restrictive at all. It is believed that the automorphism group of every vertex-transitive digraph contains a semiregular element (see [34, Problem 2.4]) and, more generally, that every 2-closed transitive permutation group contains a semiregular element [3, 24]. Substantial evidence supporting these claims has been gathered thus far. For example, it is known that all 2-closed quasiprimitive groups [16] and all 2-closed transitive groups of square-free degree [8] have this property. (See also [4, 37] for further results on this problem.)

Of course, the existence of a mixer in a bicirculant means that the two orbits of the semiregular cyclic subgroup do not form an imprimitivity block system of the full automorphism group. On the other hand, the existence of a swap in a bicirculant is equivalent to there being a transitive subgroup of automorphisms with the two orbits of the cyclic group in question as blocks of imprimitivity. Therefore, within a more general setting, the aim of this paper, essentially, is to discuss certain questions related to the following general problem.

Problem 1.1 *Let H be a semiregular abelian subgroup of a transitive permutation group G acting on a finite set X . Find conditions for (non)existence of G -invariant partitions and give their description in terms of orbits of H .*

Note that *G -invariant partitions* include imprimitivity block systems as well as the two trivial block systems of G . In the case when only the latter exist, that is when the group G is primitive, it is natural to ask whether G is simply or doubly transitive. (We remark that the concept of (im)primitivity has proven crucial in the investigation of symmetry properties of various combinatorial structures, see for

example [5, 6, 7, 40, 43, 45, 46, 47].) Thus, if G is primitive and H is regular, the above problem is related to an old question of Burnside regarding B -groups. (A group H is called a B -group if a primitive permutation group containing H as a regular subgroup is necessarily doubly transitive.) As a generalization let us call H an m - B -group if a primitive permutation group containing H as a semiregular subgroup with m orbits is necessarily doubly transitive. We may then pose the following problem.

Problem 1.2 *Given a positive integer m , determine the class of (abelian) m - B -groups.*

Of course, a 1- B -group is just a B -group. It is known that, by the classical results of Schur and Wielandt [49, Theorem 25.3 and 25.6], cyclic groups of composite order and dihedral groups are B -groups. Also, Li has recently classified all abelian B -groups [28]. As for 2- B -groups, for example, cyclic groups of prime order p are of this kind. Namely, in view of the classification of finite simple groups a primitive group of degree $2p$, $p > 5$ a prime, is doubly transitive. Studying existence of mixers for bicirculants is essentially the first step in determining which cyclic groups are 2- B -groups and which are not. On the other hand, knowing which cyclic groups are 2- B -groups is of vital importance in the study of arc-transitive bicirculants. We will deal with this question in a sequel to this paper.

To state our main theorem we first briefly introduce some notation and terminology. First, with a transitive action of a group G on X (having a semiregular abelian subgroup H with m orbits $X_1, \dots, X_m$) we associate a G -vertex-transitive digraph with vertex set X , that is, a digraph admitting G as subgroup of automorphisms, in short, a G -invariant digraph. Further, a certain matrix $\mathbf{S}$, called the *symbol of Γ relative to H* , is associated with a G -invariant digraph Γ (see the paragraphs following the statement of Theorem 1.3, and Subsection 3.2 for more details). Let λ be an eigenvalue of Γ and let H^* denote the *dual group of H* , that is, the group of all irreducible complex characters. Then we denote the set of all characters from H^* , for which λ is an eigenvalue of $\chi(\mathbf{S})$, by $K_{\mathbf{S},\lambda}$ (see the paragraphs following the statement of Theorem 1.3 and Subsection 3.3). Next, let W_λ denote the λ -eigenspace of Γ . It is well known that G acts on W_λ . Let $\mathcal{K}_\lambda$ denote the kernel of this action. The partition of X consisting of the orbits of $\mathcal{K}_\lambda$ is denoted by $\mathcal{B}_{\mathbf{S},\lambda}$. This partition induces a certain partition Δ_λ of the index-set $\{1, \dots, m\}$. (See the paragraphs following the statement of Theorem 1.3, and Section 4 for a more detailed description of the above concepts.)

The main result of this paper, Theorem 1.3 below, concerns two extremal cases related to $K_{\mathbf{S},\lambda}$. The first one deals with the situation when $K_{\mathbf{S},\lambda}$ generates the whole of H^* , while the second one covers the case when $K_{\mathbf{S},\lambda} = \{\chi_0\}$ consists of the principal character only.

Theorem 1.3 *Let G be a transitive permutation group on a finite set X containing an abelian semiregular subgroup H . Further, let Γ be a G -invariant digraph with symbol $\mathbf{S}$ and with an eigenvalue λ .*

- (i) *If $K_{\mathbf{S},\lambda}$ generates H^* and $\mathcal{K}_\lambda$ is not trivial, then the corresponding imprimitivity block system $\mathcal{B}_{\mathbf{S},\lambda}$ consists of blocks which intersect each orbit of H in at most one element. If in addition Δ_λ is universal, then each block of $\mathcal{B}_{\mathbf{S},\lambda}$ intersects each orbit of H in exactly one element. (In particular, the latter holds whenever H has prime number of orbits.)*
- (ii) *If λ is not the valency of Γ , and $K_{\mathbf{S},\lambda} = \{\chi_0\}$, then the corresponding imprimitivity block system $\mathcal{B}_{\mathbf{S},\lambda}$ consists of orbits of an intransitive normal subgroup of G containing H . If in addition Δ_λ is trivial, then the orbits of H form an imprimitivity block system of G . (In particular, the latter holds whenever H has prime number of orbits.)*

Theorem 1.3 is proved in Section 5, after introducing the required machinery in Sections 2 – 4. We briefly review the contents of each individual section.

First, coming back to Problem 1.1, we remark that in the special case when H is regular, a G -invariant partition can be obtained as the orbit-partition of a subgroup of H (see for example [28, Lemma 3.1]). In Section 2 we give an analogous description in the case when H is semiregular and abelian. Letting $X_1, \dots, X_m$ denote the orbits of H we show that a G -invariant partition $\mathcal{B}$ of X can be obtained by ‘glueing together’ (that is, taking the union of) certain orbits of some subgroup K of H (see Lemma 2.3). The way these orbits are glued together depends on an m -tuple $\underline{x} \in X_1 \times \dots \times X_m$ and on a certain partition Δ of the set $\{1, \dots, m\}$ (see the first paragraph of Section 2 for exact formulation). We call the ordered triple $(\underline{x}, \Delta, K)$ a G -triple corresponding to $\mathcal{B}$. In particular, to say that $(\underline{x}, \Delta_I, H)$ is a G -triple, where Δ_I is the trivial partition of $\{1, \dots, m\}$ consisting of 1-element subsets, means that the orbits of H are blocks of imprimitivity for the group G .

In Section 3, complex characters of abelian groups together with some additional linear algebra tools are introduced, to be used in Section 4 in the study of G -invariant partitions. To this end we use the following strategy. We consider linear actions of G on the eigenspaces of G -invariant digraphs. As already mentioned above, given such a digraph Γ , the orbits (on X) of the kernel of such an action give rise to a G -invariant partition, denoted hereafter by $\mathcal{B}_{\mathbf{S},\lambda}$, where λ is an eigenvalue of Γ corresponding to a given eigenspace, and $\mathbf{S}$ is the so called symbol of Γ relative to H , an encoding of the adjacencies of Γ through a certain $m \times m$ ‘matrix’ whose entries are subsets of H . In turn, this allows us to describe G -triples $(\underline{x}, \Delta, K)$ corresponding to the G -invariant partition $\mathcal{B}_{\mathbf{S},\lambda}$. In particular, one can find the group K from the spectral description of Γ (see Lemma 4.2). As for Δ , however, the spectral description of Γ allows us only to specify an auxiliary partition of $\{1, \dots, m\}$ whose refinement

is the partition Δ (see Lemma 4.3). As an illustration of these concepts we end Section 4 with a thorough description of all imprimitivity block systems of the cube. Let us remark that in the case when H is regular, the eigenspaces in question have a well-known interpretation as certain subspaces of the group algebra $\mathbb{C}H^*$, where H^* denotes the group of complex characters of H . In fact, they are subspaces of the Schur algebra over H^* , which is dual to the transitivity module of G over H . Schur rings have become a powerful tool in algebraic combinatorics [11]. For example, they played a central role in solving the isomorphism problem for circulant graphs [38, 39, 41], motivated by the classical Ádám's conjecture [1], as well as in some other combinatorial problems, see for example [26].

In Section 5 we describe imprimitivity block systems in two extremal cases that can arise in the context of Problem 1.1, that is, we prove Theorem 1.3, as already mentioned above. Moreover, as a consequence of Theorem 1.3 we prove that if G is primitive, then $K_{\mathbf{S},\lambda}$ must generate the whole dual group H^* whenever λ is not the valency of Γ , see Theorem 5.1. Also, we obtain a result about non-principal characters of H when H has a prime number of orbits (Corollary 5.3) as well as a refinement in the case when H has two orbits (Theorem 5.4).

Finally, in Section 6, as an application of the theory developed here (notably, Lemma 4.3 and Corollary 5.3) we give an alternative proof of an old result classifying edge-transitive generalized Petersen graphs [15] (see Theorem 6.1). In general, theorems of this nature are usually proved either by an elementary, although technically rather involved, combinatorial approach [15, 20, 32, 36, 42], or else by using a normal subgroup reduction approach, often involving the classification of finite simple groups as an essential ingredient, see for example [9, 10, 12, 13, 14, 19, 27, 28, 29, 30, 35, 44]. Our proof of Theorem 6.1 uses the second approach. However, being based on the techniques of this paper, it does not rely on the classification of finite simple groups.

2 Describing G -invariant partitions via G -triples

We start by introducing a class of partitions of a given finite set X arising from a semiregular action of a permutation group H on X . Letting $X_1, \dots, X_m$ denote the orbits of H , choose an element $\underline{x} = (x_1, \dots, x_m) \in X_1 \times \dots \times X_m$, a partition Δ of $\{1, \dots, m\}$, and a subgroup K of H . We define a partition depending on $\underline{x}$, Δ and K as follows. For each $i \in \{1, \dots, m\}$ the K -orbits contained in X_i are of the form x_i^{hK} for some $h \in H$. Moreover, since H is semiregular we have $|x_i^{hK}| = |K|$, and $x_i^{hK} = x_i^{h'K}$ if and only if $hK = h'K$. The ordered triple $(\underline{x}, \Delta, K)$ gives rise to the partition

$$\Pi(\underline{x}, \Delta, K) = \left\{ \bigcup_{i \in T} x_i^{hK} \mid T \in \Delta, h \in H \right\} \quad (1)$$

of X . We will refer to $\underline{x}$ as its *base vector*. The example below illustrates this definition.

Example 2.1 Let $X = \{1, \dots, 12\}$ and $H = \langle h \rangle$ where $h = (1\ 2\ 3\ 4)(5\ 6\ 7\ 8)(9\ 10\ 11\ 12)$. The orbits of H are $X_1 = \{1, 2, 3, 4\}$, $X_2 = \{5, 6, 7, 8\}$ and $X_3 = \{9, 10, 11, 12\}$. Choose $\Delta = \{\{1, 2\}, \{3\}\}$ and $K = \langle h^2 \rangle$. Then there are two different partitions of X of the form $\Pi(\underline{x}, \Delta, K)$, $\underline{x} \in X_1 \times X_2 \times X_3$:

base vector $\underline{x}$	elements of $\Pi(\underline{x}, \Delta, K)$
$(1, 5, 9)$	$\{1, 3, 5, 7\}, \{2, 4, 6, 8\}, \{9, 11\}, \{10, 12\}$
$(1, 6, 9)$	$\{1, 3, 6, 8\}, \{2, 4, 5, 7\}, \{9, 11\}, \{10, 12\}$

Observe that $\Pi(\underline{x}, \Delta, K) = \Pi(\underline{x}', \Delta', K')$ implies $\Delta = \Delta'$ and also $K = K'$. However, it is possible that $\underline{x} \neq \underline{x}'$. We use $\Pi_{\Delta, K}$ to denote the set of all partitions Π of X such that $\Pi = \Pi(\underline{x}', \Delta, K)$ for some $\underline{x}' \in X_1 \times \dots \times X_m$. Regarding all base vectors of a fixed partition from $\Pi_{\Delta, K}$ we have the following simple property.

Lemma 2.2 *Let H be a semiregular permutation group on a finite set X with orbits X_i , $i \in \{1, \dots, m\}$. With notation above, let $\Pi \in \Pi_{\Delta, K}$ be a partition of X . Then, given any point $x \in X$, there exists a vector $\underline{x} = (x_1, \dots, x_m)$ such that $x \in \{x_1, \dots, x_m\}$ and $\Pi = \Pi(\underline{x}, \Delta, K)$.*

PROOF. Let $\underline{y} = (y_1, \dots, y_m)$ be a base vector of $\Pi = \Pi(\underline{y}, \Delta, K)$. Further, let j be the unique index such that $x \in X_j$, and let $T \in \Delta$ such that $j \in T$. Now $x = y_j^h$ for some $h \in H$. Define the m -tuple $\underline{x} = (x_1, \dots, x_m)$ by $x_i = y_i^h$ if $i \in T$, and $x_i = y_i$ otherwise. It is easy to check that $\Pi(\underline{x}, \Delta, K) = \Pi$. ■

We denote by Δ_I the trivial partition $\{\{1\}, \dots, \{m\}\}$ of $\{1, \dots, m\}$ corresponding to the identity relation on the set $\{1, \dots, m\}$. Similarly, we let Δ_U denote the partition $\{\{1, \dots, m\}\}$ of $\{1, \dots, m\}$ corresponding to the universal relation on the set $\{1, \dots, m\}$. Obviously, $\Pi(\underline{x}, \Delta_I, K)$ does not depend on its base vector $\underline{x}$ since it is equal to the partition induced by the set of all K -orbits in X . A partition of $\{1, \dots, m\}$ is *uniform* if it consists of subsets of equal cardinality.

Let us now assume that a semiregular permutation group H on X is abelian and contained in a transitive permutation group G . In the next lemma we show that every G -invariant partition of X must then be of the form $\Pi(\underline{x}, \Delta, K)$ for some uniform partition Δ of $\{1, \dots, m\}$ and some subgroup K of H .

Lemma 2.3 *Let G be a transitive permutation group on a finite set X , and let H be a semiregular abelian subgroup of G with orbits X_i , $i \in \{1, \dots, m\}$. Let $\mathcal{B}$ be a G -invariant partition of X , let $\mathcal{K}$ be the kernel of the action of G on $\mathcal{B}$, and let*

$x \in X$. Then there exist some $\underline{x} = (x_1, \dots, x_m) \in X_1 \times \dots \times X_m$ and a uniform partition Δ of $\{1, \dots, m\}$ such that

$$\mathcal{B} = \Pi(\underline{x}, \Delta, H \cap \mathcal{K}).$$

Moreover, $\underline{x}$ can be chosen in such a way that $x \in \{x_1, \dots, x_m\}$.

PROOF. Let $B_1, B_2 \in \mathcal{B}$ and let $i \in \{1, \dots, m\}$ be such that $B_1 \cap X_i \neq \emptyset$ and $B_2 \cap X_i \neq \emptyset$. We first show that if $B_1 \cap X_j \neq \emptyset$ for some $j \in \{1, \dots, m\}$ then also $B_2 \cap X_j \neq \emptyset$. Indeed, let $x_1 \in B_1 \cap X_i$ and $x_2 \in B_2 \cap X_i$. Since X_i is an orbit of H , there is $h \in H$ such that $x_2 = x_1^h$. As $\mathcal{B}$ is a G -invariant partition of X we have $B_1^h = B_2$. But X_j is also an orbit for H ; hence $B_1 \cap X_j \neq \emptyset$ implies $B_1^h \cap X_j^h \neq \emptyset$, that is, $B_2 \cap X_j \neq \emptyset$.

Define a relation $\sim$ on the set $\{1, \dots, m\}$ by letting $i \sim j$ if and only if there exists some $B \in \mathcal{B}$ such that $B \cap X_i \neq \emptyset$ and $B \cap X_j \neq \emptyset$. Clearly, $\sim$ is a reflexive and symmetric relation. Moreover, by the arguments given in the previous paragraph, $\sim$ is also a transitive relation, and hence an equivalence relation. Let Δ denote the partition of $\{1, \dots, m\}$ induced by the equivalence classes of $\sim$. For $T \in \Delta$, set

$$\mathcal{B}_T = \{ B \in \mathcal{B} \mid B \cap X_i \neq \emptyset \text{ for all } i \in T \}.$$

Fix $T \in \Delta$ and fix a block $B_T \in \mathcal{B}_T$. Furthermore, for every $i \in T$ choose $x_i \in B_T \cap X_i$ and define K_i to be the setwise stabilizer of the set $B_T \cap X_i$ in H . Clearly, since B_T is a block and since X_i is an H -orbit, we have $K_i = K_j$ for $i, j \in T$. We may therefore define $K_T = K_i$ for $i \in T$. In fact, $K_T = H \cap \text{Stab}_{B_T}$. Moreover, $B_T = \cup_{i \in T} x_i^{K_T}$.

Now let B' be an arbitrary block in $\mathcal{B}_T$ and let $i \in T$. Since X_i is an H -orbit, there exists $h \in H$ such that $B' = B_T^h$. Therefore, since H is abelian,

$$B' = \bigcup_{i \in T} x_i^{hK_T}.$$

We show that $K_T = H \cap \mathcal{K}$, thus making K_T independent of the particular choice of $T \in \Delta$. Let $h \in H \cap \mathcal{K}$. Then $x_i^h \in B_T \cap X_i$ for every $i \in T$, implying $H \cap \mathcal{K} \subseteq K_T$. Furthermore, the action of $H\mathcal{K}/\mathcal{K}$ on $\mathcal{B}_T$ is transitive. Since $H\mathcal{K}/\mathcal{K} \cong H/(H \cap \mathcal{K})$, we have that the group $H\mathcal{K}/\mathcal{K}$ is abelian and hence regular on $\mathcal{B}_T$. It follows that $|H \cap \mathcal{K}| = |H|/|\mathcal{B}_T| = |K_T|$, and so $K_T = H \cap \mathcal{K}$ for every $T \in \Delta$.

Since T was arbitrary, we conclude that $\mathcal{B} = \Pi((x_1, \dots, x_m), \Delta, H \cap \mathcal{K})$, as required. Further, $|T| = |B_T|/|H \cap \mathcal{K}|$ for each $T \in \Delta$, and so Δ is a uniform partition. Finally, the last assertion of Lemma 2.3 follows directly from Lemma 2.2. ■

We conclude this section with the following definition. Let $\mathcal{B}$ be an arbitrary G -invariant partition of X . If $\mathcal{B} = \Pi(\underline{x}, \Delta, K)$ for some $\underline{x} \in X_1 \times \cdots \times X_m$, some uniform partition Δ of $\{1, \dots, m\}$, and a subgroup K of H , then the triple $(\underline{x}, \Delta, K)$ will be called a *G-triple corresponding to $\mathcal{B}$* .

3 G -invariant digraphs and their eigenspaces

Let G be a transitive permutation group on a finite set X . An orbit of G acting on $X \times X$ is called an *orbital* of G . To any collection of orbitals of G we can associate a vertex-transitive digraph admitting a transitive action of G , whose vertex set is X and whose edge set is induced by this collection of orbitals. Note that any digraph admitting a transitive action of G on its vertex set is necessarily of this form. As mentioned in the Introduction, we will refer to such digraphs as to *G-invariant digraphs* over X . In this section we describe, following a number of steps, the eigenspaces of G -invariant digraphs in the case when G has an abelian semiregular subgroup.

3.1 Block matrices and tensor products

In this subsection we determine the eigenvalues and eigenvectors of certain block matrices. To begin with, let us recall the definition of the tensor product. For column vectors $\mathbf{u} = (u_1, \dots, u_m)^\top \in \mathbb{C}^m$ and $\mathbf{v} = (v_1, \dots, v_n)^\top \in \mathbb{C}^n$, their *tensor product* is defined to be the mn -dimensional vector

$$\mathbf{u} \otimes \mathbf{v} = (u_1 v_1, \dots, u_1 v_n, \dots, u_m v_1, \dots, u_m v_n)^\top.$$

Let $U \leq \mathbb{C}^m$ and $W \leq \mathbb{C}^n$ be subspaces, with their respective bases $\mathbf{u}_1, \dots, \mathbf{u}_k$ and $\mathbf{w}_1, \dots, \mathbf{w}_l$. Then the *tensor product* $U \otimes W$ of U and W is the kl -dimensional subspace of $\mathbb{C}^{mn}$, spanned by vectors $\mathbf{u}_i \otimes \mathbf{w}_j$, $i \in \{1, \dots, k\}$, $j \in \{1, \dots, l\}$.

In what follows we present some technical assumptions needed in Lemma 3.1 below. Let m and n be positive integers and let

$$B = \begin{pmatrix} B_{11} & \cdots & B_{1m} \\ \vdots & \ddots & \vdots \\ B_{m1} & \cdots & B_{mm} \end{pmatrix}$$

be an $mn \times mn$ block-matrix over $\mathbb{C}$ where all blocks B_{ij} are of size $n \times n$. Note that B can be written as

$$B = \sum_{i,j=1}^m E_{ij} \otimes B_{ij},$$

where E_{ij} ($1 \leq i, j \leq m$) is the $m \times m$ matrix with the (i, j) -th entry equal to 1 and all other entries equal to 0. Suppose now that all the blocks B_{ij} have a

common orthogonal eigenvector basis consisting of (unit) column vectors $\mathbf{v}_1, \dots, \mathbf{v}_n$. For integers $i, j \in \{1, \dots, m\}$ and $k \in \{1, \dots, n\}$, let $\mu_{ij}^{(k)}$ denote the eigenvalue of B_{ij} corresponding to $\mathbf{v}_k$, and let

$$M^{(k)} = \sum_{i,j=1}^m \mu_{ij}^{(k)} E_{ij}$$

be the $m \times m$ matrix whose (i, j) -th entry is equal to $\mu_{ij}^{(k)}$. For an eigenvalue λ of $M^{(k)}$, let $V_{\lambda,k}$ denote the eigenspace of $M^{(k)}$ corresponding to λ . We denote the $m \times m$ identity matrix by I_m , and by

$$K_{B,\lambda} = \{k \mid 1 \leq k \leq n, \det(M^{(k)} - \lambda I_m) = 0\}$$

the set of all indices k such that λ is an eigenvalue of $M^{(k)}$.

Lemma 3.1 *With assumptions and notation above the following holds.*

(i) *The spectrum $\text{Spec}(B)$ is equal to the union of spectra $\cup_{k=1}^n \text{Spec}(M^{(k)})$.*

(ii) *The λ -eigenspace of B is equal to the direct sum $\bigoplus_{k \in K_{B,\lambda}} V_{\lambda,k} \otimes \langle \mathbf{v}_k \rangle$.*

PROOF. With respect to the common eigenbasis $\mathbf{v}_1, \dots, \mathbf{v}_n$, the matrix B_{ij} takes the diagonal form $D_{ij} = \text{diag}(\mu_{ij}^{(1)}, \dots, \mu_{ij}^{(n)})$, that is, $B_{ij} = P D_{ij} P^{-1}$ where $P = [\mathbf{v}_1, \dots, \mathbf{v}_n]$ is the $n \times n$ matrix with the above eigenvectors as columns. Consequently, $B = (I_m \otimes P)(\sum_{i,j=1}^m E_{ij} \otimes D_{ij})(I_m \otimes P)^{-1}$. Observe that there is a permutation matrix which establishes a similarity relation between $\sum_{i,j=1}^m E_{ij} \otimes D_{ij}$ and the block-diagonal matrix

$$\sum_{k=1}^n E_{kk} \otimes M^{(k)}.$$

It is now obvious that the spectrum of B is equal to union of spectra of all matrices $M^{(k)}$, proving (i).

In order to show (ii), let λ be an eigenvalue of B , and note that the λ -eigenspace of $\sum_{k=1}^n E_{kk} \otimes M^{(k)}$ is equal to the direct sum $\bigoplus_{k \in K_{B,\lambda}} \langle \mathbf{e}_k \rangle \otimes V_{\lambda,k}$, where $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ is the standard basis of $\mathbb{C}^n$. Since there is an automorphism of $\mathbb{C}^{mn}$ mapping the eigenspaces of $\sum_{k=1}^n E_{kk} \otimes M^{(k)}$ bijectively onto the eigenspaces of B , the dimension of the λ -eigenspace of B is equal to

$$\sum_{k \in K_{B,\lambda}} \dim V_{\lambda,k}.$$

To complete the proof we need to find an appropriate basis of this subspace. Let $\mathbf{u} \in V_{\lambda,k}$. Then $\mathbf{u} \otimes \mathbf{v}_k$ is a λ -eigenvector of B . Indeed, $B(\mathbf{u} \otimes \mathbf{v}_k) = \sum_{i,j=1}^m (E_{ij} \otimes B_{ij})(\mathbf{u} \otimes \mathbf{v}_k) = \sum_{i,j=1}^m (E_{ij} \mathbf{u} \otimes B_{ij} \mathbf{v}_k) = \sum_{i,j=1}^m (E_{ij} \mathbf{u} \otimes \mu_{ij}^{(k)} \mathbf{v}_k) = (\sum_{i,j=1}^m \mu_{ij}^{(k)} E_{ij}) \mathbf{u} \otimes \mathbf{v}_k = M^{(k)} \mathbf{u} \otimes \mathbf{v}_k = \lambda(\mathbf{u} \otimes \mathbf{v}_k)$. In each eigenspace $V_{\lambda,k}$ choose an orthogonal basis $\mathfrak{u}_{\lambda,k} = \{\mathbf{u}_{\lambda,1}^{(k)}, \dots, \mathbf{u}_{\lambda,r_k}^{(k)}\}$. Consider now two distinct eigenvectors $\mathbf{u}_{\lambda,s}^{(k)} \otimes \mathbf{v}_k$ and $\mathbf{u}_{\lambda',s'}^{(l)} \otimes \mathbf{v}_l$ of B . Their dot product is

$$\langle \mathbf{u}_{\lambda,s}^{(k)} \otimes \mathbf{v}_k, \mathbf{u}_{\lambda',s'}^{(l)} \otimes \mathbf{v}_l \rangle = \langle \mathbf{u}_{\lambda,s}^{(k)}, \mathbf{u}_{\lambda',s'}^{(l)} \rangle \langle \mathbf{v}_k, \mathbf{v}_l \rangle.$$

If $\lambda = \lambda'$, then either $k \neq l$ or else $s \neq s'$; hence this dot product is 0. Thus, $\{\mathbf{u} \otimes \mathbf{v}_k \mid k \in K_{B,\lambda}, \mathbf{u} \in \mathfrak{u}_{\lambda,k}\}$ is an orthogonal, and hence linearly independent, subset in the λ -eigenspace of the matrix B . As there are exactly $\sum_{k \in K_{B,\lambda}} \dim V_{\lambda,k}$ elements in this subset, it is actually a basis, and (ii) follows. $\blacksquare$

3.2 Symbols of G -invariant digraphs

We start by introducing some notation. Let Z be a nonempty finite set with a fixed linear ordering of its elements. By V_Z and Mat_Z we denote, respectively, the $|Z|$ -dimensional $\mathbb{C}$ -vector space of column vectors over $\mathbb{C}$, and the corresponding algebra of $|Z| \times |Z|$ matrices, both with rows (and columns) indexed by the chosen ordering of Z . (Similarly, by $\text{Mat}_Z(\mathfrak{H})$ we denote the set of $|Z| \times |Z|$ ‘matrices’ with entries in a set $\mathfrak{H}$.) It will be to our advantage to occasionally consider a complex valued function $f: Z \rightarrow \mathbb{C}$ as a vector $\mathbf{v}_f \in V_Z$, whose z -th component, $z \in Z$, is equal to $f(z)$.

Recall that for a subset S of a group H , the *Cayley digraph* $\text{Cay}(H, S)$ is the digraph with vertex set H , where there is an arc pointing from $x \in H$ to $y \in H$ whenever $yx^{-1} \in S$. (Note that this definition is more general than the usually accepted one; in particular, we allow S to contain the trivial element.) Now choose a linear ordering $h_1, h_2, \dots, h_n$ of elements of H . By $A(\text{Cay}(H, S)) \in \text{Mat}_H$ we denote the *adjacency matrix* of $\text{Cay}(H, S)$ whose (h_i, h_j) -th entry is 1 if $h_j h_i^{-1} \in S$, and is 0 otherwise.

Let G be a transitive permutation group on a finite set X and let $H \leq G$ be a semiregular subgroup. Choose a fixed linear ordering $h_1, h_2, \dots, h_n$ of elements of H and a linear ordering $X_1, \dots, X_m$ of the set $\mathfrak{X}$ of H -orbits, along with a chosen base vector $\underline{x} = (x_1, \dots, x_m)$, where $x_i \in X_i$ for each i . This naturally induces a linear ordering of X , namely

$$x_1^{h_1}, \dots, x_1^{h_n}, x_2^{h_1}, \dots, x_2^{h_n}, \dots, x_m^{h_1}, \dots, x_m^{h_n}.$$

Observe that the above is nothing else but the usual lexicographic ordering. In particular, relative to the above orderings of $\mathfrak{X}$, H , and X , we have

$$V_{\mathfrak{X}} \otimes V_H = V_X. \quad (2)$$

With assumptions and notation above, let Γ be a G -invariant digraph on X . For $x, y \in X$, we shall write $x \rightarrow_\Gamma y$ if there is an arc pointing from x to y . The $m \times m$ ‘matrix’ $\mathbf{S} \in \text{Mat}_{\mathfrak{x}}(2^H)$, whose (X_i, X_j) -th entry is the subset

$$S_{ij} = \{h \in H \mid x_i \rightarrow_\Gamma x_j^h\},$$

is called the *symbol of Γ relative to H* , the ordering of $\mathfrak{x}$, and the base vector $\underline{x}$. We use $\mathbf{S}$ to construct an adjacency matrix of Γ .

Lemma 3.2 *Let Γ be a G -invariant digraph on a finite set X , and let $H \leq G$ be a semiregular subgroup. With respect to the linear ordering of X induced by those of H and $\mathfrak{x}$ as above, the matrix $A_{\mathbf{S}} \in \text{Mat}_X$ defined by*

$$A_{\mathbf{S}} = \begin{pmatrix} A(\text{Cay}(H, S_{11})) & \cdots & A(\text{Cay}(H, S_{1m})) \\ \vdots & \ddots & \vdots \\ A(\text{Cay}(H, S_{m1})) & \cdots & A(\text{Cay}(H, S_{mm})) \end{pmatrix}$$

is an adjacency matrix of Γ .

PROOF. Pick integers $i, j \in \{1, \dots, m\}$ and $k, l \in \{1, \dots, n\}$. To prove that $A_{\mathbf{S}}$ is indeed the adjacency matrix of Γ we need to show that $x_i^{h_k} \rightarrow_\Gamma x_j^{h_l}$ if and only if there is an arc from h_k to h_l in $\text{Cay}(H, S_{ij})$. Since Γ is G -invariant, we have that $x_i^{h_k} \rightarrow_\Gamma x_j^{h_l}$ if and only if $x_i \rightarrow_\Gamma (x_j^{h_l})^{h_k^{-1}} = x_j^{h_l h_k^{-1}}$, which is equivalent to $h_l h_k^{-1} \in S_{ij}$. The result follows. $\blacksquare$

3.3 Eigenspaces of G -invariant digraphs

Let us first recall some well-known facts about eigenvalues and eigenvectors of abelian Cayley digraphs. For a finite abelian group H , let $H^* = \{\chi \mid \chi: H \rightarrow \mathbb{C}^*\}$ denote its *dual group* consisting of all irreducible complex characters (with pointwise multiplication as the group operation). For a subset $S \subseteq H$ and a character $\chi \in H^*$ we set

$$\chi(S) = \begin{cases} \sum_{g \in S} \chi(g), & S \neq \emptyset \\ 0 & S = \emptyset. \end{cases}$$

Fix a linear ordering of H . It is well-known (see for example [17, Lemma 9.2]) that the spectrum of the adjacency matrix $A(\text{Cay}(H, S)) \in \text{Mat}_H$ of the Cayley digraph $\text{Cay}(H, S)$ is the set $\{\chi(S) \mid \chi \in H^*\}$. The eigenvalue $\chi(S)$ corresponds to the eigenvector $\mathbf{v}_\chi$, and moreover, all eigenvectors $\mathbf{v}_\chi, \chi \in H^*$, form an orthogonal basis of V_H . Note that this basis does not depend on S .

The above results can be stated in a more general setting. Let G be a transitive permutation group on a finite set X having a semiregular abelian subgroup $H \leq G$ with m orbits. Fix a linear ordering of $H, \mathfrak{x}$, and X as in the previous subsection.

Further, let Γ be a G -invariant digraph and let $\mathbf{S}$ be the symbol of Γ . For a character $\chi \in H^*$ we define the $m \times m$ matrix $\chi(\mathbf{S}) \in \text{Mat}_{\mathfrak{x}}$ over $\mathbb{C}$ as follows

$$\chi(\mathbf{S}) = \begin{pmatrix} \chi(S_{11}) & \cdots & \chi(S_{1m}) \\ \vdots & \ddots & \vdots \\ \chi(S_{m1}) & \cdots & \chi(S_{mm}) \end{pmatrix}.$$

Corollary 3.3 *Let G be a transitive permutation group on a finite set X and let H be a semiregular abelian subgroup of G . Then the spectrum of a G -invariant digraph Γ on X is equal to the union of spectra of all $\chi(\mathbf{S})$, $\chi \in H^*$, where $\mathbf{S}$ is a symbol of Γ relative to H .*

PROOF. By Lemma 3.2, the spectrum of Γ is equal to the spectrum of $A_{\mathbf{S}}$. Now all of its blocks $A(\text{Cay}(H, S_{ij}))$ have a common orthogonal eigenvector basis formed by vectors $\mathbf{v}_{\chi}$, where $\chi \in H^*$, and the corresponding eigenvalue of the block $A(\text{Cay}(H, S_{ij}))$ is equal to $\chi(S_{ij})$. Thus, the matrix associated with χ is $M^{(\chi)} = \chi(\mathbf{S})$. The result then follows by applying Lemma 3.1(i) to the matrix $A_{\mathbf{S}}$. ■

Let $\text{val}(\Gamma)$ denote the *valency*, that is, the number of *out-neighbours* (or *in-neighbours*), of a G -invariant digraph Γ with symbol $\mathbf{S}$ relative to a semiregular abelian subgroup $H \leq G$. Then $\text{val}(\Gamma)$ is an eigenvalue of $\chi_0(\mathbf{S})$ (and therefore of Γ), where $\chi_0 \in H^*$ is the principal character of H .

As for the eigenvectors of $A_{\mathbf{S}}$, let λ be an eigenvalue and let $W_{\lambda} \leq V_X$ denote the respective eigenspace. First, define the set

$$K_{\mathbf{S}, \lambda} = \{ \chi \in H^* : \det(\chi(\mathbf{S}) - \lambda I_m) = 0 \} \quad (3)$$

of all characters $\chi \in H^*$ for which λ is an eigenvalue of $M^{(\chi)} = \chi(\mathbf{S})$. For $\chi \in K_{\mathbf{S}, \lambda}$, let $V_{\lambda, \chi}$ be the λ -eigenspace of the matrix $\chi(\mathbf{S})$. In view of Corollary 3.3, as a direct consequence of Lemma 3.1(ii) we obtain that W_{λ} admits the following description.

Corollary 3.4 *Let G be a transitive permutation group on a finite set X and let H be a semiregular abelian subgroup of G . Further, let Γ be a G -invariant digraph, and let $\mathbf{S}$ and $A_{\mathbf{S}}$ be its symbol and the adjacency matrix relative to H , respectively. Then with notation and assumptions above, if λ is an eigenvalue of Γ , the respective λ -eigenspace W_{λ} of $A_{\mathbf{S}}$ can be described as*

$$W_{\lambda} = \bigoplus_{\chi \in K_{\mathbf{S}, \lambda}} V_{\lambda, \chi} \otimes \langle \mathbf{v}_{\chi} \rangle.$$

4 G -invariant partitions induced by linear actions

Throughout this section, let G be a transitive permutation group on a finite set X , with $H \leq G$ an abelian semiregular subgroup having m orbits $\mathfrak{x} = \{X_1, \dots, X_m\}$, and let Γ be a G -invariant digraph on X . For fixed orderings of H , $\mathfrak{x}$, and the induced lexicographic ordering of X , let $\mathbf{S}$ be the symbol of Γ relative to H and a fixed base vector $\underline{x} = (x_1, \dots, x_m) \in X_1 \times \dots \times X_m$, and let $A_{\mathbf{S}}$ be the corresponding adjacency matrix of Γ . For an eigenvalue λ of Γ and $\chi \in K_{\mathbf{S}, \lambda}$, we let $V_{\lambda, \chi}$ and W_{λ} be the associated eigenspaces, as defined in the previous section.

It is well-known that the action of G on X has a linear extension to V_X : for an arbitrary vector $\mathbf{v} = (v_x)$, the x -th component of $\mathbf{v}^g$ is the $x^{g^{-1}}$ -th component of $\mathbf{v}$, that is, G acts by the rule $(v_x)^g = (v_{xg})$. The restriction of this action to H is described in the next lemma.

Lemma 4.1 *With notation and assumptions above, let $\mathbf{u} \in V_{\mathfrak{x}}$ and let $f: H \rightarrow \mathbb{C}$ be an arbitrary function. Then for $h \in H$ we have*

$$(\mathbf{u} \otimes \mathbf{v}_f)^h = \mathbf{u} \otimes \mathbf{v}_{f_h}, \quad (4)$$

where $f_h: H \rightarrow \mathbb{C}$ is defined by the rule $f_h(a) = f(ah)$.

PROOF. Write $\mathbf{u} \otimes \mathbf{v}_f = (v_x)$, $x \in X$. In view of (2) we have $v_x = u_i f(a)$ whenever $x = x_i^a$. Similarly, letting $\mathbf{u} \otimes \mathbf{v}_{f_h} = (w_x)$ we have $w_x = u_i f_h(a) = u_i f(ah)$ whenever $x = x_i^a$. Note that if $x = x_i^a$, then $x^h = x_i^{ah}$, implying $v_{x^h} = u_i f(ah) = w_x$. Hence $(\mathbf{u} \otimes \mathbf{v}_f)^h = (v_x)^h = (v_{x^h}) = (w_x) = \mathbf{u} \otimes \mathbf{v}_{f_h}$. $\blacksquare$

For $g \in G$, let $P_g \in \text{Mat}_X$ denote the permutation matrix representing the action of g on X . Since Γ is G -invariant, P_g commutes with $A_{\mathbf{S}}$. Hence if λ is an eigenvalue of $A_{\mathbf{S}}$, then the corresponding eigenspace W_{λ} is an invariant subspace for the linear action of G on V_X . Let K_{λ} be the kernel of this action of G on W_{λ} , and let $\mathcal{B}_{\mathbf{S}, \lambda}$ denote the partition of X consisting of the orbits of K_{λ} . Note that for each $\mathbf{w} \in W_{\lambda}$ the coordinates of $\mathbf{w}$, indexed by the elements in the same orbit of K_{λ} , are the same. Clearly, $\mathcal{B}_{\mathbf{S}, \lambda}$ is a G -invariant partition, and so by Lemma 2.3 we have that

$$\mathcal{B}_{\mathbf{S}, \lambda} = \Pi(\underline{y}, \Delta_{\lambda}, K_{\lambda} \cap H), \quad (5)$$

for a suitable base vector $\underline{y} = (y_i) \in X_1 \times \dots \times X_m$ and a uniform partition Δ_{λ} of $\{1, \dots, m\}$, with K_{λ} denoting the kernel of the action of G on $\mathcal{B}_{\mathbf{S}, \lambda}$. Obviously, $K_{\lambda} \leq K_{\lambda}$. In fact, $K_{\lambda} = K_{\lambda}$. To show this, let $w = (w_x) \in W_{\lambda}$ and $g \in K_{\lambda}$. Since $w^g = (w_{xg})$, it suffices to see that $w_x = w_{xg}$ for $x \in X$. But this is obvious because if $x, y \in X$ are in the same orbit of K_{λ} , then $w_x = w_y$ (by definition of K_{λ}). Our goal in this section is to examine the G -triple given in (5).

First we consider the group $\mathcal{K}_\lambda \cap H$ as in (5). To this end, the following correspondence $\perp$ (also known as the *Dirichlet correspondence*) between the set of subgroups of H and the set of subgroups of H^* will play an essential role. For $L \leq H$ and $L^* \leq H^*$, we let

$$\begin{aligned} L^\perp &= \{ \chi \in H^* : L \leq \text{Ker} \chi \} \\ L^{*\perp} &= \{ h \in H : \chi(h) = 1 \text{ for all } \chi \in L^* \} = \cap_{\chi \in L^*} \text{Ker} \chi. \end{aligned}$$

Lemma 4.2 *With assumptions and notation above we have $\mathcal{K}_\lambda \cap H = \langle K_{\mathbf{S}, \lambda} \rangle^\perp$.*

PROOF. Recall that $\mathcal{K}_\lambda = K_\lambda$ is the kernel of the linear action of G on the eigenspace W_λ . By Corollary 3.4 it follows that $g \in \mathcal{K}_\lambda$ if and only if

$$(\mathbf{u} \otimes \mathbf{v}_\chi)^g = \mathbf{u} \otimes \mathbf{v}_\chi$$

for each $\chi \in K_{\mathbf{S}, \lambda}$ and for each $\mathbf{u} \in V_{\lambda, \chi}$. Applying Lemma 4.1 for $u \in V_\chi$, $h \in H$ and $\chi \in H^*$, we obtain

$$(\mathbf{u} \otimes \mathbf{v}_\chi)^h = \mathbf{u} \otimes (\chi(h) \mathbf{v}_\chi) = \chi(h)(\mathbf{u} \otimes \mathbf{v}_\chi).$$

Now let $h \in \mathcal{K}_\lambda \cap H$. Then, combining the above two facts together we get that $\chi(h) = 1$ for all $\chi \in K_{\mathbf{S}, \lambda}$. Hence $\mathcal{K}_\lambda \cap H \leq \langle K_{\mathbf{S}, \lambda} \rangle^\perp$. Conversely, take $h \in \langle K_{\mathbf{S}, \lambda} \rangle^\perp \leq H$. Then, by definition of $\langle K_{\mathbf{S}, \lambda} \rangle^\perp$ we have $\chi(h) = 1$ for all $\chi \in K_{\mathbf{S}, \lambda}$. Again, combining the above facts it follows that $w^h = w$ for each $w \in W_\lambda$, and hence $h \in \mathcal{K}_\lambda$. This completes the proof of Lemma 4.2. $\blacksquare$

We now examine the partition Δ_λ of (5). Since Δ_λ is not readily at hand, we introduce certain auxiliary partitions of $\{1, \dots, m\}$ which are easier to compute, and, as it will become clear shortly, they all have Δ_λ as a refinement. Hence their intersection gives at least some information about Δ_λ , the more so when the former is close to the identity partition.

Let λ be an eigenvalue of Γ and let $\chi \in K_{\mathbf{S}, \lambda}$. Choose a basis $\mathfrak{u}$ of $V_{\lambda, \chi}$. For $k \in \{1, \dots, m\}$ we let $\mathfrak{u}^{(k)}$ denote the vector formed by the k -th components of all vectors in $\mathfrak{u}$. The auxiliary partition $\Delta_{\lambda, \chi}$ is now defined in terms of the following equivalence relation $\sim_{\lambda, \chi}$ on $\{1, \dots, m\}$:

$$i \sim_{\lambda, \chi} j \iff \mathfrak{u}^{(i)} = \chi(h) \mathfrak{u}^{(j)} \text{ for some } h \in H.$$

Note that, given λ and χ , the partition $\Delta_{\lambda, \chi}$ does not depend on the chosen basis $\mathfrak{u}$. Given two arbitrary partitions Π_1 and Π_2 of $\{1, \dots, m\}$ we write $\Pi_1 \sqsubseteq \Pi_2$ to indicate that Π_1 is a refinement of Π_2 .

Lemma 4.3 *With assumptions and notation above, for all λ and $\chi \in K_{\mathbf{S},\lambda}$ we have that $\Delta_\lambda \subseteq \Delta_{\lambda,\chi}$.*

PROOF. Fix a character $\chi \in K_{\mathbf{S},\lambda}$ and a basis $\mathfrak{u}$ of $V_{\lambda,\chi}$. Let i and j be in the same partition class of Δ_λ . We are going to show that $i \sim_{\lambda,\chi} j$. Recalling that $(x_1, \dots, x_m)$ is the base vector of the symbol $\mathbf{S}$, consider $x_i \in X_i$ and $x_j \in X_j$. By Lemma 2.3, the orbit of $\mathcal{K}_\lambda$ containing x_i intersects also X_j . Hence there exists $h \in H$ such that x_i and x_j^h are in the same $\mathcal{K}_\lambda$ -orbit. Consider now a base vector $\mathbf{u} = (u_1, u_2, \dots, u_m)^\top \in \mathfrak{u}$. By Corollary 3.4, $\mathbf{u} \otimes \mathbf{v}_\chi \in W_\lambda$. Since x_i and x_j^h are in the same $\mathcal{K}_\lambda$ -orbit, the x_i -th and x_j^h -th coordinate of $\mathbf{u} \otimes \mathbf{v}_\chi$ are the same. Thus, in view of (2) we have $u_i = \chi(h)u_j$, and since $\mathbf{u} \in \mathfrak{u}$ was arbitrary, $\mathfrak{u}^{(i)} = \chi(h)\mathfrak{u}^{(j)}$. In other words, $i \sim_{\lambda,\chi} j$. $\blacksquare$

As an illustration of the above tools we analyse the cube Q_3 with $G \leq \text{Aut}(Q_3)$ an arbitrary vertex-transitive subgroup. Note that there are total of 8 possible choices (conjugate classes) for G : $\text{Aut}(Q_3) \cong S_4 \times \mathbb{Z}_2$ acting 2-arc-transitively on Q_3 , two conjugate classes of groups of order 24 acting 1-arc-transitively on Q_3 , one group of order 16 and four groups of order 8 (acting regularly on Q_3), two of which are isomorphic to dihedral group of order 8, and the other two isomorphic to $\mathbb{Z}_4 \times \mathbb{Z}_2$ and $\mathbb{Z}_2^3$, respectively.

Note that the cube has two essentially different representations as a bicirculant: $[S_{11}, S_{22}, S_{12}] = [\{1, 3\}, \{1, 3\}, \{0\}]$ and $[S_{11}, S_{22}, S_{12}] = [\emptyset, \emptyset, \{1, 2, 3\}]$. In both cases H is isomorphic to $\mathbb{Z}_4$. The orbits are denoted by X_1 and X_2 , and (x_1, x_2) is the base vector. To simplify the notation we identify H with $\mathbb{Z}_4$. In particular, we may write $X_1 = \{x_1^0, x_1^1, x_1^2, x_1^3\}$ and $X_2 = \{x_2^0, x_2^1, x_2^2, x_2^3\}$. The first example gives a thorough analysis for the case $[S_{11}, S_{22}, S_{12}] = [\{1, 3\}, \{1, 3\}, \{0\}]$, while in the second case $[S_{11}, S_{22}, S_{12}] = [\emptyset, \emptyset, \{1, 2, 3\}]$ some of the technical details are omitted.

Example 4.4 Let $[S_{11}, S_{22}, S_{12}] = [\{1, 3\}, \{1, 3\}, \{0\}]$. Then

$$\mathbf{S} = \begin{pmatrix} \{1, 3\} & \{0\} \\ \{0\} & \{1, 3\} \end{pmatrix}.$$

Therefore, the neighbors of x_i^j are $x_i^{j\pm 1}$ and x_{i+1}^j , where $i \in \mathbb{Z}_2$ and $j \in \mathbb{Z}_4$. We first compute the eigenvalues of Q_3 via the matrices $\chi(\mathbf{S})$, $\chi \in H^* = \{\chi_0, \chi_1, \chi_2, \chi_3\}$, where $\chi_j(h) = \mathbf{i}^{jh}$ for $j \in \{0, 1, 2, 3\}$ and $h \in \mathbb{Z}_4$, with $\mathbf{i}$ denoting the imaginary unit. It transpires that the eigenvalues of $\chi_0(\mathbf{S})$ are 1 and 3, the eigenvalues of $\chi_1(\mathbf{S})$ and $\chi_3(\mathbf{S})$ are 1 and -1 , and that the eigenvalues of $\chi_2(\mathbf{S})$ are -1 and -3 . Therefore,

$$K_{\mathbf{S},3} = \{\chi_0\}, \quad K_{\mathbf{S},1} = \{\chi_0, \chi_1, \chi_3\}, \quad K_{\mathbf{S},-1} = \{\chi_1, \chi_2, \chi_3\}, \quad K_{\mathbf{S},-3} = \{\chi_2\}.$$

Using Lemma 4.2 we find that

$$\mathcal{K}_3 \cap H = H, \quad \mathcal{K}_{-3} \cap H = \{0, 2\}, \quad \mathcal{K}_1 \cap H = \mathcal{K}_{-1} \cap H = \{0\}.$$

Furthermore, the subspaces $V_{\lambda, \chi}$, $\lambda \in \{3, 1, -1, -3\}$, $\chi \in K_{\mathbf{S}, \lambda}$, are the following ones:

$$V_{\lambda, \chi} = \begin{cases} \langle (1, 1)^\top \rangle & (\lambda, \chi) \in \{(3, \chi_0), (1, \chi_1), (1, \chi_3), (-1, \chi_2)\} \\ \langle (1, -1)^\top \rangle & (\lambda, \chi) \in \{(1, \chi_0), (-1, \chi_1), (-1, \chi_3), (-3, \chi_2)\}. \end{cases}$$

Therefore, by the definition of the partition $\Delta_{\lambda, \chi}$, $\lambda \in \{3, 1, -1, -3\}$, $\chi \in K_{\mathbf{S}, \lambda}$, we have that Δ_{1, χ_0} is the trivial partition $\{\{1\}, \{2\}\}$, and that $\Delta_{\lambda, \chi}$ is the universal partition $\{\{1, 2\}\}$ in all other cases. By Lemma 4.3 we obtain that Δ_1 is the trivial partition, while in all other cases Δ_λ is either the trivial or the universal partition. Since $\mathcal{K}_1 \cap H = \{0\}$ and Δ_1 is the trivial partition, the corresponding block system $\mathcal{B}_{\mathbf{S}, 1}$ given in (5) is the trivial block system consisting of singletons. In all other cases, direct analysis of the action of the group G on the corresponding eigenspace W_λ is needed to compute Δ_λ , and consequently, the block system $\mathcal{B}_{\mathbf{S}, \lambda}$.

Consider first the case $\lambda = 3$. Since $W_3 = \langle (1, 1, 1, 1, 1, 1, 1)^\top \rangle$, we obtain that $\mathcal{K}_3 = G$, and so Δ_3 is universal since G is vertex-transitive. Therefore, the corresponding block system $\mathcal{B}_{\mathbf{S}, 3}$ is trivial with the whole set of vertices as the only block.

Next, consider the case $\lambda = -3$ and observe $W_{-3} = \langle (1, -1, 1, -1, -1, 1, -1)^\top \rangle$. Clearly, $\mathcal{K}_{-3}$ is the largest subgroup in G which fixes the bipartition $\{\{x_1^0, x_1^2, x_2^1, x_2^3\}, \{x_1^1, x_1^3, x_2^0, x_2^2\}\}$. Note that irrespective what a group G is, the group $\mathcal{K}_{-3}$ has to be of index 2 in G , and therefore acts transitively on each of the two sets of bipartition. Hence, Δ_{-3} is the universal partition, and consequently the corresponding block system $\mathcal{B}_{\mathbf{S}, -3}$ coincides with the bipartition of Q_3 .

Finally, consider the case $\lambda = -1$ and observe that

$$W_{-1} = \langle (1, -1, 1, -1, 1, -1, 1)^\top, (1, \mathbf{i}, -1, -\mathbf{i}, -1, -\mathbf{i}, 1)^\top, (1, -\mathbf{i}, -1, \mathbf{i}, -1, \mathbf{i}, 1)^\top \rangle.$$

In this case $\mathcal{K}_{-1}$ is the largest subgroup in G fixing the antipodal partition $\{\{x_1^0, x_2^2\}, \{x_1^1, x_2^3\}, \{x_1^2, x_2^0\}, \{x_1^3, x_2^1\}\}$. By viewing Q_3 as the canonical double cover of the complete graph K_4 with the fibres corresponding to the pairs of antipodal vertices, it may be seen that $\mathcal{K}_{-1}$ is transitive on the pairs of antipodal vertices precisely when G is a lift of a transitive subgroup of $\text{Aut}(K_4) \cong S_4$, that is when G is isomorphic to one of the following groups: $S_4 \times \mathbb{Z}_2$, $A_4 \times \mathbb{Z}_2$, $D_8 \times \mathbb{Z}_2$, $\mathbb{Z}_4 \times \mathbb{Z}_2$ or $\mathbb{Z}_2^3$. In these five cases the partition Δ_{-1} is the universal partition, and the corresponding block system $\mathcal{B}_{\mathbf{S}, -1}$ coincides with the antipodal partition of Q_3 . The remaining three possibilities for the group G give rise to trivial partition Δ_{-1} , and therefore to the trivial block system $\mathcal{B}_{\mathbf{S}, -1}$ consisting of singletons.

Example 4.5 Let $[S_{11}, S_{22}, S_{12}] = [\emptyset, \emptyset, \{1, 2, 3\}]$. Then

$$K_{\mathbf{S},3} = K_{\mathbf{S},-3} = \{\chi_0\}, \quad K_{\mathbf{S},1} = K_{\mathbf{S},-1} = \{\chi_1, \chi_2, \chi_3\}.$$

Using Lemma 4.2 we find that

$$\mathcal{K}_3 \cap H = \mathcal{K}_{-3} \cap H = H, \quad \mathcal{K}_1 \cap H = \mathcal{K}_{-1} \cap H = \{0\}.$$

Similarly as in the previous example we find that Δ_{-3,χ_0} is the trivial partition, while $\Delta_{\lambda,\chi}$ is the universal partition in all other cases. Therefore, Δ_{-3} is the trivial partition, and the corresponding block system $\mathcal{B}_{\mathbf{S},-3}$ coincides with the bipartition of Q_3 . In all other cases, a direct analysis of the action of the group G on the corresponding eigenspace W_λ is needed to compute Δ_λ , and consequently, the block system $\mathcal{B}_{\mathbf{S},\lambda}$. Below are the conclusions.

If $\lambda = 3$ then the corresponding block system $\mathcal{B}_{\mathbf{S},3}$ is trivial with the whole set of vertices as the only block.

If $\lambda = 1$ then (by considering an arbitrary basis for W_1) it is easy to see that $\mathcal{K}_1$ is trivial. Therefore Δ_1 is the trivial partition and gives rise to the trivial block system $\mathcal{B}_{\mathbf{S},3}$ consisting of singletons.

Finally, consider the case $\lambda = -1$. By considering an arbitrary basis for W_{-1} we find that $\mathcal{K}_{-1}$ is the largest subgroup in G fixing the antipodal partition

$$\{\{x_1^0, x_2^0\}, \{x_1^1, x_2^1\}, \{x_1^2, x_2^2\}, \{x_1^3, x_2^3\}\}.$$

Hence, as in the previous case, Δ_{-1} is the universal partition (and the corresponding block system $\mathcal{B}_{\mathbf{S},-1}$ coincides with the antipodal partition of Q_3) if G is isomorphic to $S_4 \times \mathbb{Z}_2$, $A_4 \times \mathbb{Z}_2$, $D_8 \times \mathbb{Z}_2$, $\mathbb{Z}_4 \times \mathbb{Z}_2$, and $\mathbb{Z}_2^3$. In the remaining three possibilities for the group G , $\mathcal{B}_{\mathbf{S},-1}$ is the trivial block system consisting of singletons.

In each of these two examples a complete list of block systems for the full automorphism group (but also of some transitive subgroups) is obtained. However, there are additional block systems in Q_3 arising from some vertex-transitive but not arc-transitive subgroups of $\text{Aut}Q_3$. To obtain these block systems, a modification of the above approach is needed. For example, by letting the group G in Example 4.4 be the regular group isomorphic to $\mathbb{Z}_4 \times \mathbb{Z}_2$ (or the appropriate regular dihedral group D_8), and by replacing the whole eigenspace W_1 with the subspace generated by the vector $(1, 1, 1, 1, -1, -1, -1, -1)^\top$, we obtain the block system consisting of the orbits of the subgroup H (isomorphic to $\mathbb{Z}_4$).

5 Analyzing (im)primitivity block systems

5.1 Existence of (im)primitivity block systems

With the results of Section 4 we are now ready to address the main topic of this paper regarding imprimitivity block systems of transitive permutation groups containing abelian semiregular subgroups. Following discussion in Examples 4.4 and 4.5, two types of block systems seem to stand out. First, when $\mathcal{K}_\lambda \cap H = 1$ with Δ_λ nontrivial, and second, when $\mathcal{K}_\lambda \cap H = H$ with Δ_λ not universal.

The antipodal block systems in Examples 4.4 and 4.5 fall within a framework of a general situation when $\mathcal{K}_\lambda \cap H = 1$ with Δ_λ nontrivial, implying that $\mathcal{B}_{\mathbf{S},\lambda}$ is an imprimitivity block system consisting of blocks which intersect each orbit of H in at most one element. This is considered in Theorem 1.3(i). The bipartition block system in Example 4.5 falls within a framework of a general situation when $\mathcal{K}_\lambda \cap H = H$ with Δ_λ not universal, implying that $\mathcal{B}_{\mathbf{S},\lambda}$ is an imprimitivity block system consisting of unions of orbits of H ; see Theorem 1.3(ii). We are now ready to prove Theorem 1.3.

Proof of Theorem 1.3. First, observe that in view of the fact that $K_{\mathbf{S},\lambda}$ generates H^* , we have $\mathcal{K}_\lambda \cap H = 1$ by Lemma 4.2. Consequently, each block of $\mathcal{B}_{\mathbf{S},\lambda}$ intersects each orbit of H in at most one element. Second, since $\mathcal{K}_\lambda$ is not trivial, the partition Δ_λ is nontrivial, implying that $\mathcal{B}_{\mathbf{S},\lambda}$ is an imprimitivity block system.

The special case when Δ_λ is universal is now obvious. In particular, since the partition is uniform and not trivial, Δ_λ must indeed be universal whenever H has prime number of orbits. This proves (i).

In order to prove (ii), observe that in view of the fact that $K_{\mathbf{S},\lambda} = \{\chi_0\}$, we have $\mathcal{K}_\lambda \cap H = H$ by Lemma 4.2. Second, since $\lambda \neq \text{val}(\Gamma)$ we have that $(1, 1, \dots, 1)^\top \notin V_{\lambda, \chi_0}$. Therefore, Δ_{λ, χ_0} is not universal by the definition of the relation $\sim_{\lambda, \chi_0}$. By Lemma 4.3, also the partition Δ_λ is not universal. The desired intransitive normal subgroup of G is $\mathcal{K}_\lambda$.

The special case when Δ_λ is trivial is now obvious. In particular, since Δ_λ is uniform and not universal, the partition must indeed be trivial whenever H has prime number of orbits. ■

Besides these two extreme situations there are other interesting block systems whose structure can be easily described. For example, if the number of H -orbits is prime and $\lambda \neq \text{val}(\Gamma)$, then the orbits of $\langle K_{\mathbf{S},\lambda} \rangle^\perp$ obviously form a G -invariant partition of X . Furthermore, using Theorem 1.3(ii), the group $\langle K_{\mathbf{S},\lambda} \rangle$ can be fully described in the special case when G is a primitive group.

Theorem 5.1 *Let G be a primitive permutation group containing an abelian semiregular subgroup H and let Γ be a G -invariant digraph with symbol $\mathbf{S}$. If $\lambda \neq \text{val}(\Gamma)$ is an eigenvalue of Γ , then $\langle K_{\mathbf{S},\lambda} \rangle = H^*$.*

PROOF. As G is primitive, $\mathcal{B}_{\mathbf{S},\lambda}$ consists either of singletons, or else comprises the whole set X . In the first case we have $\mathcal{K}_\lambda \cap H = 1$ (with Δ_λ trivial), implying $\langle K_{\mathbf{S},\lambda} \rangle^\perp = 1$, that is, $\langle K_{\mathbf{S},\lambda} \rangle = H^*$. In the second case we have $\mathcal{K}_\lambda \cap H = H$ (with Δ_λ universal), implying $\langle K_{\mathbf{S},\lambda} \rangle^\perp = H$, that is, $K_{\mathbf{S},\lambda} = \{\chi_0\}$. Since $\lambda \neq \text{val}(\Gamma)$, Theorem 1.3(ii) implies that G is imprimitive, a contradiction. ■

An immediate consequence of Theorem 5.1 is the following result dealing with primitive groups containing regular abelian subgroups. Based on this result, a method for investigation of B-groups was developed by Burnside [2]. As an application it was proved that cyclic p -groups of composite order are B-groups for any prime p , see also [25].

Corollary 5.2 *Let G be a primitive permutation group containing an abelian regular subgroup H , and let $\Gamma = \text{Cay}(H, S)$ be a G -invariant Cayley digraph with symbol $\mathbf{S} = [S]$. If $\lambda \neq \text{val}(\Gamma)$ is an eigenvalue of Γ , then $\langle K_{\mathbf{S},\lambda} \rangle = \langle \{\chi \in H^* \mid \chi(S) = \lambda\} \rangle = H^*$.*

5.2 Mixers and bi-Cayley graphs

Let G be a transitive permutation group on a finite set X containing an abelian semiregular subgroup H . Generalizing the notion of a mixer of a bicirculant introduced in Section 1, we say that $g \in G$ is a *mixer relative to H* (in short, a *mixer* when the subgroup H is clear from the context), if the orbits of H are not blocks of imprimitivity for $\langle g \rangle$. The following corollary is just a rephrasing of a special case of Theorem 1.3(ii) into the mixers language.

Corollary 5.3 *Let G be a transitive permutation group on a finite set X containing an abelian semiregular subgroup H with a prime number of orbits, and let Γ be a G -invariant digraph with symbol $\mathbf{S}$. If G has a mixer relative to H , then for any eigenvalue $\lambda \neq \text{val}(\Gamma)$ such that $\chi_0 \in K_{\mathbf{S},\lambda}$ we have $K_{\mathbf{S},\lambda} \neq \{\chi_0\}$.*

When the number of orbits of H is relatively small, some additional information about mixers can be obtained using a more direct combinatorial approach which takes into account the intersections of orbits of H with their images under a mixer. We here consider the case of two orbits.

Let G be a transitive permutation group on X and H a semiregular abelian subgroup having two orbits X_1 and X_2 . Fix a base vector $(x_1, x_2) \in X_1 \times X_2$, and

let Γ be a G -invariant digraph (in this context also known as the *bi-Cayley digraph*) with symbol $\mathbf{S}$ relative to the chosen base vector. Note that

$$\mathbf{S} = \begin{pmatrix} S & T \\ Q & R \end{pmatrix},$$

with $|S| = |R|$ and $|T| = |Q|$. Therefore, the eigenvalues of $\chi_0(\mathbf{S})$ are $\text{val}(\Gamma) = |S| + |T|$ and $d(\Gamma) = |S| - |T|$.

For $g \in G$ we define the subsets M_g and N_g of H by

$$\begin{aligned} M_g &= \{h \in H \mid (x_1^h)^g \in X_1\}, \\ N_g &= \{h \in H \mid (x_2^h)^g \in X_1\}. \end{aligned}$$

Since g is an automorphism we have $|M_g| + |N_g| = |H|$. Observe that g is a swap if and only if $M_g = \emptyset$ and $N_g = H$, and that g is a mixer if and only if $M_g \neq \emptyset$ and $M_g \neq H$.

In what follows it will be convenient to view subsets of H as elements of the group algebra $\mathbb{Q}H$. Following [49], a subset A of H is considered as $\underline{A} = \sum_{h \in A} h \in \mathbb{Q}H$. Characters of H are naturally extended to an algebra-homomorphism $\mathbb{Q}H \rightarrow \mathbb{C}$ by letting $\chi(\alpha) = \sum_{h \in H} a_h \chi(h)$ for $\alpha = \sum_{h \in H} a_h h$.

Theorem 5.4 *Let G be a transitive permutation group containing an abelian semiregular subgroup H having two orbits, and let Γ be a G -invariant digraph. Then with notation and assumptions above, for all $g \in G$ and for all characters $\chi \in H^* \setminus K_{\mathbf{S}, d(\Gamma)}$ we have*

$$\chi(M_g) = \chi(N_g) = 0.$$

PROOF. Let $g \in G$. For convenience we use a shorthand notation $M = M_g$ and $N = N_g$. Choose $h \in H$. The number of out-neighbours of x_1^h that are contained in $M \cup N$ is equal to $|hS \cap M| + |hT \cap N|$. This number is equal to the coefficient of h in $\underline{S}^{-1} \underline{M} + \underline{T}^{-1} \underline{N} \in \mathbb{Q}H$. Moreover, observe that it is also equal to the number of out-neighbours of x_1^{hg} that are contained in X_1 , which is either $|S|$ or $|Q|$ ($= |T|$), depending on whether $h \in M$ or $h \in H \setminus M$. Consequently,

$$\underline{S}^{-1} \underline{M} + \underline{T}^{-1} \underline{N} = |S| \underline{M} + |T| \underline{H \setminus M} = d(\Gamma) \underline{M} + |T| \underline{H}.$$

Similarly, by counting the number of out-neighbours of x_2^h that are contained in $M \cup N$ (which equals the number of out-neighbours of x_2^{hg} that are contained in X_1) we obtain the equation

$$\underline{Q}^{-1} \underline{M} + \underline{R}^{-1} \underline{N} = |S| \underline{N} + |T| \underline{H \setminus N} = d(\Gamma) \underline{N} + |T| \underline{H}.$$

Applying a nonprincipal character $\chi \in H^*$ to both of the above equalities we find that $\chi(M)$ and $\chi(N)$ are solutions of the following system of linear equations

$$\begin{pmatrix} (\chi(S^{-1}) - d(\Gamma)) & \chi(T^{-1}) \\ \chi(Q^{-1}) & (\chi(R^{-1}) - d(\Gamma)) \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

If $\chi \notin K_{\mathbf{S}, d(\Gamma)}$ we have $\det(\chi(\mathbf{S}) - d(\Gamma)I) = (\chi(S) - d(\Gamma))(\chi(R) - d(\Gamma)) - \chi(T)\chi(Q) \neq 0$. As the determinant of the above system of linear equations is the conjugate of $\det(\chi(\mathbf{S}) - d(\Gamma)I)$, the system has only trivial solution. Thus, $\chi(M) = \chi(N) = 0$. $\blacksquare$

We remark that in the particular case when H has two orbits, Corollary 5.3 follows directly from Theorem 5.4. Namely, let $g \in G$ be a mixer relative to H . Then M_g and N_g are nonempty proper subsets of H . It is well known that if all non-principal characters vanish on a subset $A \subseteq H$, then either $A = \emptyset$ or $A = H$; see for instance [48, Corollary 1.3.5]. Therefore, there exists a non-principal character $\chi \in H^*$ such that either $\chi(M_g) \neq 0$ or $\chi(N_g) \neq 0$. By Theorem 5.4, we have $\chi \in K_{\mathbf{S}, d(\Gamma)}$.

6 An application: generalized Petersen graphs

To further illustrate the possible use of techniques developed here we give an alternative short proof of a well know classical result about edge-transitivity (and thus arc-transitivity) of generalized Petersen graphs due to Frucht, Graver and Watkins [15, Theorem 2]. Other applications will appear elsewhere.

Theorem 6.1 *A generalized Petersen graph $\Gamma = \text{GP}(n, s)$ is edge-transitive (and thus arc-transitive) if and only if (n, s) is one of the following pairs: $(4, 1)$, $(5, 2)$, $(8, 3)$, $(10, 2)$, $(10, 3)$, $(12, 5)$, or $(24, 5)$.*

Recall that the *generalized Petersen graph* $\text{GP}(n, s)$ is the bicirculant relative to the cyclic group $H = \mathbb{Z}_n$ and symbol

$$\mathbf{S} = \begin{pmatrix} \{1, -1\} & \{0\} \\ \{0\} & \{s, -s\} \end{pmatrix},$$

where $s \in \mathbb{Z}_n \setminus \{0\}$.

As remarked in Section 1, theorems of this nature are usually proved either by an elementary, although technically rather involved, combinatorial approach (as, for instance, counting the number of 8-cycles in the original proof of the above theorem), or else by using a two-step (normal subgroup) reduction approach, where the first

step identifies a small restricted family from which all other graphs in the class can be reconstructed, and the second step consists in the actual construction of these graphs. Our proof of Theorem 6.1 uses the above two steps approach. The first step is based on the techniques of this paper, more precisely, on Corollary 5.3 and Lemma 4.2. The second step uses two simple observations about covers of the cube and of the Petersen graph.

Proof of Theorem 6.1. We split the proof into two parts. In the first part we show that every edge-transitive generalized Petersen graph Γ is a regular cyclic cover either of the cube or of the Petersen graph (such that $\text{Aut}(\Gamma)$ projects). In the second part we determine all such covers.

1. PART. First note that the eigenvalues of $\chi_0(\mathbf{S})$ are $\text{val}(\Gamma) = 3$ and $d(\Gamma) = 1$. The respective eigenspace V_{1,χ_0} is spanned by $(1, -1)^\top$. By the definition of the relation $\sim_{1,\chi_0}$ the partition Δ_{1,χ_0} is trivial, and by Lemma 4.3 the partition Δ_1 is also trivial. It follows that $\mathcal{K}_1 \leq H$. Thus, the orbits of $L = \mathcal{K}_1 \cap H = \mathcal{K}_1$ are contained in the orbits of H . Since the L -orbits form an $\text{Aut}(\Gamma)$ -invariant partition of the vertex set of Γ , the group $\text{Aut}(\Gamma)$ projects along the cyclic regular covering projection $\Gamma \rightarrow \Gamma/L$. In order to determine L and consequently Γ/L , recall that, by Lemma 4.2 and by the Dirichlet correspondence, $L = \langle K_{\mathbf{S},1} \rangle^\perp = \cap \text{Ker} \chi$, where χ ranges over all characters in $K_{\mathbf{S},1}$. Therefore, we first find $K_{\mathbf{S},1}$.

The set $K_{\mathbf{S},1}$ is determined using the assumption that Γ is edge-transitive, and hence that it has a mixer relative to H . By Corollary 5.3, the set $K_{\mathbf{S},1}$ contains a non-principal character, say χ , implying that

$$(\chi(1) + \chi(-1) - 1)(\chi(s) + \chi(-s) - 1) = 1. \quad (6)$$

All solutions of (6) give all possible characters in $K_{\mathbf{S},1}$. (Observe that $\chi \in K_{\mathbf{S},1}$ if and only if $\chi^{-1} \in K_{\mathbf{S},1}$.) For a natural number m let $\xi_m = e^{\frac{2\pi i}{m}}$ denote the ‘smallest’ primitive m -th root of unity. As $\chi(1) = \xi_n^j$ for some $j \in \mathbb{Z}_n \setminus \{0\}$, equation (6) rewrites as $(\xi_n^j + \xi_n^{-j} - 1)(\xi_n^{js} + \xi_n^{-js} - 1) = 1$. By setting $j' = j/\gcd(j, n)$, $n' = n/\gcd(j, n)$, and $s' = s \bmod n'$, this last equation becomes

$$(\xi_{n'}^{j'} + \xi_{n'}^{-j'} - 1)(\xi_{n'}^{j's'} + \xi_{n'}^{-j's'} - 1) = 1.$$

Since j' and n' are coprime, $\xi_{n'}^{j'}$ and $\xi_{n'}$ are both primitive n' -th roots of unity. Hence their minimal polynomial is the same, implying that

$$(\xi_{n'} + \xi_{n'}^{-1} - 1)(\xi_{n'}^{s'} + \xi_{n'}^{-s'} - 1) = 1. \quad (7)$$

If both factors in (7) are positive then $n' = 1$, forcing $j = n$. However, this is not possible as χ is non-principal. Therefore $\xi_{n'} + \xi_{n'}^{-1} - 1 = 2 \cos(2\pi/n') - 1 < 0$, and so $n' \leq 5$. A direct analysis yields that either $n' = 4$ and $s' = \pm 1$, or else

$n' = 5$ and $s' = \pm 2$. It follows that $j' = \pm 1$ in the first case while $j' \in \{\pm 1, \pm 2\}$ in the second case. Hence for some natural number m either $n = 4m$, $j = \pm m$, and $s = \pm 1 \pmod{4}$, or else $n = 5m$, $j \in \{\pm m, \pm 2m\}$, and $s = \pm 2 \pmod{5}$. We now analyze each of these cases separately.

Let $n = 4m$, $j = \pm m$, and $s = \pm 1 \pmod{4}$, where m is not divisible by 5. Since $\chi(1) = \xi_n^j \in \{\xi_4, \xi_4^{-1}\}$ we have that $K_{\mathbf{S},1} = \{\chi_0, \chi, \chi^{-1}\}$, and so $L = \langle K_{\mathbf{S},1} \rangle^\perp$ is the unique index 4 subgroup in $\mathbb{Z}_n$. Since $s = \pm 1 \pmod{4}$, the graph Γ/L is the cube.

Let $n = 5m$, $j \in \{\pm m, \pm 2m\}$, and $s = \pm 2 \pmod{5}$ where m is not divisible by 4. Since $\chi(1) = \xi_n^j \in \{\xi_5, \xi_5^{-1}, \xi_5^2, \xi_5^{-2}\}$ we have that $K_{\mathbf{S},1} = \{\chi_0, \chi, \chi^{-1}, \chi^2, \chi^{-2}\}$, and so $L = \langle K_{\mathbf{S},1} \rangle^\perp$ is the unique index 5 subgroup in $\mathbb{Z}_n$. Since $s = \pm 2 \pmod{5}$, the graph Γ/L is the Petersen graph.

Let $n = 20m$ where $s = \pm 1 \pmod{4}$ and $s = \pm 2 \pmod{5}$. By the above analysis, the group $\langle K_{\mathbf{S},1} \rangle = \langle \chi, \chi' \rangle$ where $\chi(1) = \xi_4$ and $\chi'(1) = \xi_5$. In this case $L = \langle K_{\mathbf{S},1} \rangle^\perp$ is the unique index 20 subgroup in $\mathbb{Z}_n$. Since $s = \pm 1 \pmod{4}$ and $s = \pm 2 \pmod{5}$ we have, by the Chinese Remainder Theorem, that $s = \pm 3 \pmod{20}$ or $s = \pm 7 \pmod{20}$. Hence Γ/L is either $\text{GP}(20, 3)$ or $\text{GP}(20, 7)$. In particular, Γ/L and hence also Γ is a regular cyclic cover of both the cube and the Petersen graph. (Actually, as it will become clear in Part 2, this case cannot occur.)

Note that in all of the above cases L is isomorphic to $\mathbb{Z}_m$.

2. PART. Recall from Part 1 that $\Gamma \rightarrow \Gamma/L$ is a regular cyclic covering projection such that $\text{Aut}(\Gamma)$ projects, and hence some minimal arc transitive subgroup of Γ/L lifts. We determine all such covers where the base graph is either the cube or the Petersen graph. Since $\text{GP}(20, 3)$ or $\text{GP}(20, 7)$ are not among them, the case where the base graph is one of these two need not be considered.

As it is well known, see for instance [18], any regular $\mathbb{Z}_m$ -covering projection $\tilde{Y} \rightarrow Y$ can be reconstructed by means of voltages $\zeta(i, j) \in \mathbb{Z}_m$ assigned to arcs (i, j) of Y , with inverse arcs carrying inverse values; moreover, without loss of generality one can assume that the arcs of some arbitrarily chosen spanning tree in Y are assigned the trivial voltage 0. In order to determine whether a particular automorphism $\alpha \in \text{Aut}(Y)$ lifts along $\tilde{Y} \rightarrow Y$ we use the theory developed in [33]. In our case, the induced action of α on base homology cycles, must extend to an automorphism of $\mathbb{Z}_m$.

Let Y be the cube. Denote its vertices by $\{0, 1, 2, 3, 4, 5, 6, 7\}$ in such a way that 01230 and 45674 are the outer and the inner cycles, respectively, and 04, 15, 26 and 37 are the spokes. Choose a spanning tree consisting of edges 01, 12, 23, 04, 15, 26, and 37. The base homology cycles are $C_1 = 30123$, $C_2 = 45104$, $C_3 = 56215$, $C_4 = 67326$, and $C_5 = 7401237$. Voltages on the spanning tree are 0. Without loss of generality we can assume that the voltages on the remaining arcs are $\zeta(3, 0) = 1$, and $\zeta(4, 5) = \zeta(5, 6) = \zeta(6, 7) = a$, and $\zeta(7, 4) = a + 1$. (This is obtained using the fact that the derived covering graph is the generalized Petersen graph.) Consider now the automorphism $\alpha = (134)(527)$ which must lift since an edge-transitive group of

Γ projects. Since the induced mapping of voltages of $\zeta(C_j) \rightarrow \zeta(C_j^\alpha)$ must extend to an automorphism $x \mapsto \lambda x$ of L , we get the following system of equations in L :

$$\lambda = a, \quad \lambda a = 1, \quad \lambda a = a, \quad \lambda a = -4a - 1, \quad \lambda(a + 1) = 2a.$$

This gives $\lambda = a = 1$ and $6 = 0$ in L , implying that L is isomorphic to $\mathbb{Z}_2$, $\mathbb{Z}_3$, or $\mathbb{Z}_6$. If $L = \mathbb{Z}_2$ then $\Gamma \cong \text{GP}(8, 5) \cong \text{GP}(8, 3)$. If $L = \mathbb{Z}_3$ then $\Gamma \cong \text{GP}(12, 5)$. If $L = \mathbb{Z}_6$ then $\Gamma \cong \text{GP}(24, 5)$.

Let Y be the Petersen graph. Denote its vertices by $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ in such a way that 012340 and 567895 are the outer and the inner cycles, respectively, and 05, 16, 27, 38, and 49 are the spokes. Choose a spanning tree consisting of edges 01, 12, 23, 34, 05, 16, 27, 38, and 49. The base homology cycles are $C_1 = 401234$, $C_2 = 572105$, $C_3 = 683216$, $C_4 = 794327$, $C_5 = 8501238$, and $C_6 = 9612349$. Voltages on the spanning tree are 0. Without loss of generality we can assume that the voltages on the remaining arcs are $\zeta(4, 0) = 1$, and $\zeta(5, 7) = \zeta(6, 8) = \zeta(7, 9) = a$, and $\zeta(8, 5) = \zeta(9, 6) = a + 1$. (This is obtained using the fact that the derived covering graph is the generalized Petersen graph.) Consider now the automorphism $\alpha = (154)(289)(367)$ which must lift since an edge-transitive group of Γ projects. Since the induced mapping of voltages of $\zeta(C_j) \rightarrow \zeta(C_j^\alpha)$ must extend to an automorphism $x \mapsto \lambda x$ of L , we get the following system of equations in L :

$$\lambda = -2a - 1, \quad \lambda a = a, \quad \lambda a = 5a + 2, \quad \lambda a = a, \quad \lambda(a + 1) = -3a - 1, \quad \lambda(a + 1) = -3a - 1.$$

This gives $2 = 0$, $\lambda = -1$ and either $a = 0$ or $a = 1$ in L , implying that L is isomorphic to $\mathbb{Z}_2$. If $a = 0$ then $\Gamma \cong \text{GP}(10, 2)$. If $a = 1$ then $\Gamma \cong \text{GP}(10, 7) \cong \text{GP}(10, 3)$, completing the proof of Theorem 6.1. ■

References

- [1] A. Ádám, “Research problem 2-10”, *J. Combin. Theory* **2** (1967), 393.
- [2] W. Burnside, “Theory of Groups of Finite Order”, 2nd ed., Cambridge Univ. Press, Cambridge, UK, 1911.
- [3] P. J. Cameron (ed.), Problems from the Fifteenth British Combinatorial Conference, *Discrete Math.* **167/168** (1997), 605–615.
- [4] P. J. Cameron, M. Giudici, W. M. Kantor, G. A. Jones, M. H. Klin, D. Marušič, and L. A. Nowitz, Transitive permutation groups without semiregular subgroups, *J. London Math. Soc.* **6** (2002), 325–333.
- [5] E. Dobson, On solvable groups and circulant graphs, *European J. Combin.* **21** (2000), 881–885.
- [6] E. Dobson, On groups of odd prime-power degree that contain a full cycle, *Discrete Math.* **299** (2005), 65–78.
- [7] E. Dobson, Automorphism groups of metacirculant graphs of order a product of two distinct primes, *Comb. Prob. Comp.* **15** (2006), 105–130.

- [8] E. Dobson, A. Malnič, D. Marušič and L. A. Nowitz, Minimal normal subgroups of transitive permutation groups of square-free degree, *Discrete Math.*, in print.
- [9] X. G. Fang and C. E. Praeger, Finite two-arc transitive graphs admitting a Suzuki simple group, *Comm. Algebra* **27** (1999), 3727–3754.
- [10] X. G. Fang and C. E. Praeger, Finite two-arc transitive graphs admitting a Ree simple group, *Comm. Algebra* **27** (1999), 3755–3769.
- [11] I. A. Faradžev, M. H. Klin and M. Muzychuk, “Cellular rings and groups of automorphisms of graphs”, In: Investigations on Algebraic Theory of Combinatorial Objects, Math. and Its Appl. (eds. I. A. Faradžev et al.), Kluwer Acad. Publ. **84** (1994), 1–152.
- [12] Y. Q. Feng, J. H. Kwak and K. S. Wang, Classifying cubic symmetric graphs of order $8p$ or $8p^2$, *European J. Combin.* **26** (2005), 1033–1052.
- [13] Y. Q. Feng and J. H. Kwak, Classifying cubic symmetric graphs of order $10p$ or $10p^2$, *Science in China Ser A* **49**, 300–319.
- [14] Y. Q. Feng and J. H. Kwak, One-regular cubic graphs of order a small number times a prime or a prime square, *J. Austral. Math. Soc.* **76** (2004), 345–356.
- [15] R. Frucht, J. E. Graver and M. E. Watkins, The groups of the generalized Petersen graphs, *Proc. Camb. Phil. Soc.* **70** (1971), 211–218.
- [16] M. Giudici, Quasiprimitive groups with no fixed point free elements of prime order, *J. London Math. Soc.* **67** (2003), 73–84.
- [17] C. Godsil, “Algebraic Combinatorics”, Chapman & Hall, New York, 1993.
- [18] J. L. Gross and T. W. Tucker, “Topological Graph Theory”, Wiley - Interscience, New York, 1987.
- [19] A. Hassani, L. R. Nochefranca and C. E. Praeger, Two-arc transitive graphs admitting a two-dimensional projective linear group, *J. Group Theory* **4** (1999), 335–353.
- [20] J. D. Horton and I. Z. Bouwer, Symmetric Y -graphs and H -graphs, *J. Combin. Theory Ser. B* **53** (1991), 114–129.
- [21] A. A. Ivanov, Graphs of girth 5 and diagram geometries connected with the Petersen graph, *Soviet Math. Dokl.* **36** (1988), 83–87.
- [22] A. A. Ivanov, R. Liebler, T. Penttila and C. Praeger, Antipodal distance-transitive covers of complete bipartite graphs, *European J. Combin.* **18** (1997), 11–33.
- [23] A. A. Ivanov and C. Praeger, On finite affine 2-arc transitive graphs, *European J. Combin.* **14** (1993), 421–444.
- [24] M. Klin, personal communication.
- [25] W. Knapp, On Burnside’s Method, *J. Algebra* **175** (1995), 644–660.
- [26] I. Kovacs, Classifying arc transitive circulants, *J. Algebraic Combin.* **21** (2004), 353–358.
- [27] C. H. Li, The finite vertex-primitive and vertex-biprimitive s -transitive graphs for $s \geq 4$, *Trans. Amer. Math. Soc.* **353** (2001), 3511–3529.
- [28] C. H. Li, The finite primitive permutation groups containing an abelian regular subgroup, *Proc. London Math. Soc.* **87** (2003), 725–747.

- [29] C. H. Li, Permutation groups with a cyclic regular subgroup and arc transitive circulants, *J. Algebraic Combin.* **21** (2005), 131–136.
- [30] C. H. Li, Z. P. Lu and H. Zhang, Tetravalent edge-transitive Cayley graphs with odd number of vertices, *J. Combin. Theory Ser. B* **96** (2006), 164–181.
- [31] A. Malnič, Group actions, coverings and lifts of automorphisms, *Discrete Math.* **182** (1998), 203–218.
- [32] A. Malnič, D. Marušič and N. Seifter, Constructing infinite one-regular graphs, *European J. Combin.* **20** (1999), 845–853.
- [33] A. Malnič, R. Nedela and M. Škovič, Lifting graph automorphisms by voltage assignments, *European J. Combin.* **21** (2000), 927–947.
- [34] D. Marušič, On vertex symmetric digraphs, *Discrete Math.* **36** (1981), 69–81.
- [35] D. Marušič, On 2-arc-transitive Cayley graphs, *J. Combin. Theory Ser. B* **87** (2003), 162–196.
- [36] D. Marušič, Half-transitive group actions on finite graphs of valency 4, *J. Combin. Theory Ser. B* **73** (1998), 41–76.
- [37] D. Marušič and R. Scapellato, Permutation groups, vertex-transitive digraphs and semiregular automorphisms, *European J. Combin.* **19** (1998), 707–712.
- [38] M. Muzychuk, Ádám’s conjecture is true in the square-free case, *J. Combin. Theory Ser. A* **72** (1995), 118–134.
- [39] M. Muzychuk, On Ádám’s conjecture for circulant graphs, *Discrete Math.* **176** (1997), 285–298.
- [40] M. Muzychuk, On normal quotients of transitive graphs, *Ars Combinatoria* **59** (2001), 171–180.
- [41] M. Muzychuk, A solution of the isomorphism problem for circulant graphs, *Proc. London Math. Soc.* **88** (2004), 1–41.
- [42] J. M. Oh and K. W. Wang, Construction of one-regular graphs of valency 4 and 6, *Discrete Math.* **278** (2004), 195–207.
- [43] C. E. Praeger, Imprimitive symmetric graphs, *Ars Combinatoria* **19A** (1985), 149–163.
- [44] C. E. Praeger, Finite normal edge-transitive Cayley graphs, *Bull. Austral. Math. Soc.* **6** (1999), 207–220.
- [45] C. E. Praeger, Quotients and inclusions of finite quasiprimitive permutation groups, *J. Algebra* **269** (2003), 329–346.
- [46] C. E. Praeger, Quasiprimitivity: structure and combinatorial applications, *Discrete Math.* **264** (2003), 211–224.
- [47] C. E. Praeger, Finite transitive permutation groups and bipartite vertex-transitive graphs, *Illinois J. Math.* **47** (2003), 461–475.
- [48] B. Schmidt, “Characters and Cyclotomic Fields in Finite Geometry”, Lecture Note in Mathematics, Springer-Verlag, Berlin, 2002.
- [49] H. Wielandt, “Finite Permutation Groups”, Academic Press, New York 1964.